

Weakly bound nuclei and realistic NN interactions

E. N. E. van Dalen, P. Gögelein, and H. Mütter
*Institut für Theoretische Physik,
Universität Tübingen, D-72076 Tübingen, Germany*

Nuclei close to the neutron drip line are described employing an interaction model which is based on the low-momentum interaction V_{lowk} . This effective two-body interaction which is determined to reproduce the nucleon-nucleon (NN) scattering data at energies below the pion threshold is supplemented by a density-dependent contact interaction fitted to reproduce the saturation properties of infinite nuclear matter within the Hartree-Fock approach. It is demonstrated that corresponding calculations for closed shell-nuclei using this interaction model reproduce the bulk properties of these nuclei, independent whether the wave functions are expanded in terms of harmonic oscillator waves or in a basis of plane waves discretized in a spherical box of appropriate size. This plane wave basis, however, is more appropriate to describe weakly bound nuclei and the transition from discrete nuclei to homogeneous matter which is supposed to occur e.g. in the crust of neutron stars. Properties of exotic nuclei are studied within a Hartree-Fock plus BCS approximation.

PACS numbers: 21.60.Jz, 21.30.Fe, 21.65.-f, 26.60.Gj

I. INTRODUCTION

The new generation of radioactive beam facilities, e.g. the future GSI facility FAIR in Germany, the Rare Isotope accelerator planned in the United States of America or SPIRAL2 at GANIL/France, facilitate the nuclear structure studies away from the line of β -stability, especially for the neutron-rich nuclei. The study of these nuclei is of high interest since they represent important steps in the nuclear reaction chains for the formation of elements. The weak binding energies of these nuclei and the corresponding large spatial extensions of the orbits for the valence nucleons lead to very interesting features like e.g. a neutron halo made of several neutrons outside the core.

These nuclei can furthermore be understood as a first step in the transition from isolated nuclei to infinite homogeneous matter, which should occur in the outer crust of a neutron star. This transition is a very good example for a quantum liquid with a phase transition from the droplet to the homogeneous phase. This transition is a challenge for theoretical nuclear structure physics as it incorporates the transition from isolated nuclei via a crystal-like structure of quasi-nuclei embedded in a sea of neutrons to the phase of homogeneous baryon matter. These structures have been described by means of Thomas-Fermi calculations or mean-field calculations in a Wigner-Seitz cell using simple phenomenological models for the nucleon-nucleon (NN) interaction like the density-dependent Skyrme forces[1, 2, 3, 4, 5, 6] or a relativistic mean-field approach[7].

All these investigations are based on phenomenological interactions, which have been adjusted to describe the saturation properties of symmetric nuclear matter and the structure of stable nuclei with large binding energies located in the valley of β -equilibrium. These models and interactions have been developed to provide a simple description of the mean field in terms of local single-particle densities. The predictive power of such simple phenomenological interactions, however, may be rather limited as these models have been constructed to fit the data of finite nuclei.

An alternative to this phenomenological approach is the use of so-called realistic interactions like the charge-dependent Bonn interaction[8] or one of the Argonne interaction models[9]. Such realistic interactions are based e.g. on the meson exchange model and adjusted to describe the experimental data of nucleon-nucleon (NN) scattering. These interactions contain rather strong short range and tensor components, which make it inevitable to employ non-perturbative approximation schemes for the solution of the many-body problem for the nuclear hamiltonian based on such interactions[10]. Nuclear structure calculations using such interactions have become feasible for very light nuclei and provide a very good description if an appropriate three-nucleon force has been added[11]. In the next future such sophisticated calculations will not be possible for heavier nuclei or the nuclear structures in the crust of neutron stars mentioned above.

A possible way out of this problem is to consider an interaction model, which separates the low-momentum (below a cut-off Λ) and high-momentum components of a realistic NN interaction by means of renormalization techniques[12, 13, 14, 15]. If the cutoff Λ is chosen around $\Lambda = 2 \text{ fm}^{-1}$ the resulting low-momentum interaction V_{lowk} still describes the NN scattering data up to the pion threshold and turns out to be independent of the underlying realistic interaction V . Since the high-momentum components, which correspond to the short-distance behavior, of V have been removed, the resulting V_{lowk} does not produce short-range correlations in a significant way, but can be treated within the Hartree-Fock approximation[15].

Employing V_{lowk} in a calculation of nuclear matter, however, one obtains a binding energy per nucleon increasing

with density in a monotonic way, such that no saturation point is obtained[15, 16]. In order to compensate this deficiency we will supplement V_{lowk} by a simple density-dependent contact interaction adjusted to reproduce the empirical saturation property of symmetric nuclear matter. This density-dependent contact interaction, which is introduced to obtain the empirical saturation point of nuclear matter, can also be interpreted as a parameterization of the three-nucleon force, which, as discussed above, seems to be necessary for a microscopic description of nuclear structure data based on realistic NN interactions. Therefore we had the hope to obtain a good description of finite nuclei using this interaction model. We will demonstrate below that this hope has been fulfilled to a good extent.

Because of the non-locality of V_{lowk} nuclear structure calculations of finite nuclei using this interaction are typically not performed in coordinate space but employ a set of appropriate single-particle wave functions as a set of basis states, typically those of a harmonic oscillator potential. The choice of the harmonic oscillator (HO) basis seems to be rather plausible if one wants to describe the structure deeply bound, double magic nuclei like ${}^4\text{He}$, ${}^{16}\text{O}$, and ${}^{40}\text{Ca}$ [17, 18]. This choice may be questionable if nuclei close to the proton or neutron drip line are to be considered. The HO basis may not be appropriate to describe the tail of the single-particle wave functions for the weakly bound valence nucleons. The main reason, why this oscillator basis is so popular, must be attributed to the fact that the HO basis provides a simple transformation from relative coordinates in coordinate or momentum space to a coordinate system located in a laboratory frame. Such a transformation is required to allow for structure calculation of finite nuclei using realistic interaction as defined e.g. by V_{lowk} .

The HO basis is certainly not appropriate to describe the quasi-nuclear structures in the pasta phase of neutron stars mentioned above. For those studies a basis of plane wave (PW) states confined to a finite box with a size large to the nuclear dimension may be more suitable and could be considered as an appropriate basis for a Wigner-Seitz cell calculation[5, 7]. One of the main aims of this work is to establish the techniques, which are needed to allow for nuclear structure calculations of finite nuclear systems using realistic interactions in such a PW basis.

In this work we describe double magic nuclei and weakly bound neutron-rich nuclei in the framework of Hartree-Fock (HF) plus BCS calculation using a V_{lowk} potential supplemented with a contact interaction. We are going to investigate nuclei up to ${}^{208}\text{Pb}$ and compare the results of calculations using HO and PW basis states.

The plan of this paper is as follows. The procedure to renormalize the low-momentum interaction and obtain the matrix elements of V_{lowk} is reviewed in Sec. II. In this section we also describe our interaction model supplementing V_{lowk} by an appropriate contact interaction. We also work out how to solve the HF equation in a basis of harmonic oscillator and plane wave states. This section is completed by a short description of the BCS procedure. The results for infinite matter and finite nuclei are presented and discussed in Sec. III. Finally, Sec. IV contains a summary and the conclusions of our work.

II. MODEL OF THE NN INTERACTION

The idea behind the V_{lowk} interaction model is to separate the low-momentum and high-momentum components of realistic interactions and restrict the nuclear structure calculation to the low-momentum components. This means that one tries to define a model space, which accounts for the low-momentum degrees of freedom and renormalizes the effective Hamiltonian for this low-momentum regime to account for the effects of the high-momentum parts, which are integrated out. A model space technique to disentangle these parts, the unitary-model-operator approach (UMOA) [19], can be used to calculate the matrix elements for V_{lowk} .

First, we define the projection operators P , which projects onto the low-momentum subspace, and Q , which projects onto the complement of this subspace, as in all well known model space techniques. Furthermore, these operators satisfy the usual relations like $P + Q = 1$, $P^2 = P$, $Q^2 = Q$, and $PQ = 0 = QP$. The idea of the unitary-model-operator approach is now to define a unitary transformation U so that the transformed Hamiltonian does not couple P and Q , which means

$$QU^{-1}HUP = 0 \quad (1)$$

has to be fulfilled. The effective interaction is defined in terms of this unitary transformation as

$$V_{eff} = U^{-1}(h_0 + v_{12})U - h_0, \quad (2)$$

with v_{12} representing the bare NN -interaction and a starting Hamiltonian h_0 describing the one-body part of the two-body system. The unitary operator U can be written as

$$U = (1 + \omega - \omega^\dagger)(1 + \omega\omega^\dagger + \omega^\dagger\omega)^{-1/2} \quad (3)$$

with an operator ω fulfilling $\omega = Q\omega P$ and $\omega^2 = \omega^{\dagger 2} = 0$. In [15] the operator ω is obtained by first solving the two-body eigenvalue equation

$$(h_0 + v_{12})|\Phi_k\rangle = E_k|\Phi_k\rangle \quad (4)$$

and afterwards defining the matrix elements of ω using the eigenstates $|\Phi_p\rangle$ having the largest overlap with the P space. With the corresponding U then the effective interaction V_{eff} is calculated as described in [15, 20]. In this way one obtains an effective Hamiltonian $H_{eff} = h_0 + V_{eff}$, which contains this effective interaction V_{eff} . The eigenvalues, which are obtained by diagonalizing this effective Hamiltonian in the model-space (P -space), are identical to the diagonalization of the original Hamiltonian $h_0 + V$ in the complete space.

This model-space scheme can now be applied to the effective two-nucleon problem by considering for the basis states of the 2-nucleon system the states identified by the relative momentum, its modulus and the corresponding partial waves. For a given partial wave the states of the P -space are identified as those states with a relative momentum smaller than a cutoff Λ . Applying the technique described above leads to the effective interaction V_{lowk} . Solving now the Schroedinger equation for V_{lowk} within the model space, e.g. by solving the Lipmann Schwinger equation for NN scattering with the cutoff Λ , yields the same phase shift as solving the Lipmann Schwinger equation without cutoff for the original interaction v_{12} . If the original interaction v_{12} is realistic in the sense, that it has been fitted to the experimental NN phase-shifts, these phase shifts, up to the cutoff Λ , will also be reproduced by V_{lowk} .

Furthermore, the resulting V_{lowk} is found to be essentially model independent. This means that the potential is independent of the underlying realistic interaction v_{12} , if the cutoff Λ is chosen around $\Lambda = 2 \text{ fm}^{-1}$. Thus, one is able to extract a low-momentum potential V_{lowk} , which in a model independent manner describes the low-momentum component of realistic NN interactions in a more or less unique way. Note, that, by construction, this V_{lowk} interaction is defined in terms of matrix elements in a basis of NN states labelled by relative momentum.

The fact that the high-momentum or short-range components of realistic NN interactions have been integrated out by means of the unitary transformation of Eq. (3), seems to lead to problems in the description of the saturation behavior of nuclear matter. Because of this lack of short-range correlation effects, which are modified in the medium, the emergence of a saturation point is prevented in calculations of symmetric nuclear matter [13, 16]. In order to achieve saturation in symmetric nuclear matter, three-body interaction terms or density-dependent two-nucleon interactions are needed. Therefore, the effective interaction V_{lowk} is supplemented by a simple contact interaction, which we have chosen following the notation of the Skyrme interaction to be of the form

$$\Delta\mathcal{V} = \Delta\mathcal{V}_0 + \Delta\mathcal{V}_3, \quad (5)$$

with

$$\Delta\mathcal{V}_0 = \frac{1}{4}t_0[(2+x_0)\rho^2 - (2x_0+1)(\rho_n^2 + \rho_p^2)] \quad (6)$$

and

$$\Delta\mathcal{V}_3 = \frac{1}{24}t_3\rho^\alpha[(2+x_3)\rho^2 - (2x_3+1)(\rho_n^2 + \rho_p^2)], \quad (7)$$

where ρ_p and ρ_n refer to the local densities for protons and neutrons while the matter density is denoted as $\rho = \rho_p + \rho_n$. The parameters of the contact interaction are t_0 , x_0 , t_3 , x_3 and α . As described below we have chosen a fixed value of $\alpha = 0.5$ and $x_0 = 0.0$ and fitted t_0 and t_3 and x_3 in such a way that HF calculations using V_{lowk} plus the contact term of Eq. (5) yield the empirical saturation point for symmetric nuclear matter and reproduce the symmetry energy at saturation density.

Within the HF approximation this interaction model leads to a single-particle Hamiltonian for protons and neutrons ($\nu = n, p$) of the form

$$H_{HF,\nu} = H_{kin} + H_{lowk,\nu} + \Delta H_{ct,\nu} + H_{Coul}\delta_{\nu p}, \quad (8)$$

where H_{kin} is the kinetic part and $\Delta H_{ct,\nu}$ originates from the contact interaction of Eq. (5). This part is given by

$$\begin{aligned} \Delta H_{ct,\nu} = & t_0/2[(2+x_0)\rho - (1+2x_0)\rho_\nu] + t_3/24[(2+x_3)(2+\alpha)\rho^{1+\alpha} \\ & - (2x_3+1)\{2\rho^\alpha\rho_\nu + \alpha\rho^{\alpha-1}(\rho_n^2 + \rho_p^2)\}]. \end{aligned} \quad (9)$$

Furthermore, for the charged particles, the protons, one has an additional contribution to the hamiltonian, the Coulomb contribution. It is given by

$$H_{Coul} = U_{coul,dir} + U_{Coul,exch}, \quad (10)$$

where the direct term is

$$U_{coul,dir} = 4\pi e^2 \left\{ \begin{array}{ll} \int dr' r'^2 \rho_p(r')/r & \text{for } r' \leq r \\ \int dr' r' \rho_p(r') & \text{for } r' \geq r \end{array} \right. \quad (11)$$

and the exchange term is

$$U_{coul,exch} = -e^2 \left(\frac{3}{\pi} \right)^{1/3} \rho_p^{1/3}. \quad (12)$$

Note that $\Delta H_{ct,i}$ and H_{Coul} are local. They are defined in terms of the single-particle densities ρ_i resulting from the eigenstates of $H_{HF,i}$, which implies that they have to be determined in a self-consistent way, as usual for HF calculations.

As it has been mentioned above, the effective interaction V_{lowk} is non-local and defined in terms of matrix elements. This implies that the HF calculation has to be performed in a Hilbert space using an appropriate basis $|\alpha\rangle, |\beta\rangle, \dots$. The HF Hamiltonian is then expressed in terms of the matrix elements between these basis states $\langle\alpha|H_{HF}|\beta\rangle$ and the HF single-particle states $|\Psi_n\rangle$ are defined in terms of the expansion coefficients in this basis

$$|\Psi_n\rangle = \sum_{\alpha} |\alpha\rangle \langle\alpha|\Psi_n\rangle = \sum_{\alpha} c_{n\alpha} |\alpha\rangle. \quad (13)$$

The part of the HF Hamiltonian originating from V_{lowk} can be expressed in terms of two-body matrix elements by

$$\langle\alpha|H_{lowk}|\beta\rangle = \sum_{\gamma,\delta} \langle\alpha\gamma|V_{lowk}|\beta\delta\rangle \rho_{\gamma\delta}, \quad (14)$$

with $\rho_{\gamma\delta}$ representing the single-particle density matrix.

If we restrict the HF variational procedure to a spherical description of nuclei, as we will do in the following, the HF Hamiltonian is diagonal in the angular momentum quantum numbers j, l, m and the expansion is simplified to

$$\langle\vec{r}|\Psi_n\rangle = \Psi_{nljm}(\vec{r}) = \sum_{i=1}^{\infty} c_{nilj} \Phi_{iljm}(\vec{r}) \approx \sum_{i=1}^N c_{nilj} \Phi_{iljm}(\vec{r}), \quad (15)$$

where c_{nilj} are expansion coefficients and $\Phi_{iljm}(\vec{r}) = \langle\vec{r}|\alpha\rangle$ are the wave functions of our orthogonal basis. The number of basis states N has to be chosen to guarantee that the results are not affected by this limitation. Note, that using this coordinate space representation of the basis states allows the calculation of the matrix elements of $\langle\alpha|H_{HF}|\beta\rangle$ for local terms in H_{HF} in a straight forward way.

In this paper calculations are done using two different orthogonal basis systems. These basis systems are respectively that of a free particle and that of a spherical harmonic oscillator. In these cases the wave functions of our orthogonal basis can be separated in a radial part and an angular part,

$$\Phi_{iljm}(\vec{r}) = \langle\vec{r}|iljm\rangle = R_{il}(r) \mathcal{Y}_{ljm}(\vartheta, \varphi). \quad (16)$$

In both basis systems, $\mathcal{Y}_{ljm}$ represent the spherical harmonics including the spin degrees of freedom by coupling the orbital angular momentum l with the spin to a single-particle angular momentum j .

The radial part R_{il} is different for the two basis systems. The first basis system considered is that of a particle moving free in a spherical cavity with a radius R , the plane wave (PW) basis [5]. The radial wave functions R_{il} are then given by the spherical Bessel functions,

$$R_{il}(r) = N_{il} j_l(k_{il}r), \quad (17)$$

for the discrete momenta k_{il} , which fulfill

$$R_{il}(R) = N_{il} j_l(k_{il}R) = 0 \Rightarrow j_l(k_{il}R) = 0. \quad (18)$$

The normalization constant in Eq. (17) is given by

$$N_{il} = \begin{cases} i\pi \frac{\sqrt{2}}{\sqrt{R^3}} & \text{for } l = 0, \\ \frac{1}{j_{l-1}(k_{il}R)} \frac{\sqrt{2}}{\sqrt{R^3}} & \text{for } l > 0. \end{cases} \quad (19)$$

It ensures that the basis functions are orthogonal and normalized within the box,

$$\int_0^R d^3r \Phi_{iljm}^*(\vec{r}) \Phi_{i'l'j'm'}(\vec{r}) = \delta_{ii'} \delta_{ll'} \delta_{jj'} \delta_{mm'}. \quad (20)$$

Next, the basis system of the harmonic oscillator (HO) is considered. The radial part of the basis functions takes the form [21]

$$R_{il}(r) = \sqrt{\frac{2(i!)}{a^3 \Gamma(i+l+3/2)}} (r/a)^l \exp(-0.5 r^2/a^2) L_i^{l+1/2}(r^2/a^2), \quad (21)$$

where $L_i^{l+1/2}$ is a Laguerre polynomial, Γ is a gamma function, and $a = \sqrt{\hbar/(m\omega)}$ is the oscillator length. These basis functions Φ_{iljm} are orthogonal and normalized within a box with an infinite radius. However, in practice one can use a radius $R \gg a$ due to the exponential factor $\exp(-0.5 r^2/a^2)$ in the radial basis function of Eq. (21).

The matrix elements of V_{lowk} or other realistic NN interactions, which are required in Eq.(14), are easily calculated within the HO basis: In a first step one can calculate the oscillator matrix elements using the momentum representation of the relative basis. The second step, transforming these matrix elements from the relative system to the laboratory system is also straight forward using the well known transformation brackets of the Talmi-Moshinsky transformation[21, 22, 23].

While the corresponding first step, the calculation of matrix elements of V_{lowk} in the relative basis is trivial in the case of PW states, the transformation from relative to laboratory coordinates is much less convenient in the PW than in the HO basis. The evaluation and use of these vector brackets has been described e.g. in [24, 25] and will be used here.

The Hartree-Fock approximation discussed so far is a good approximation for double closed shell nuclei. In order to consider nuclei with open shells as well, we are going to use a rather phenomenological treatment of pairing correlations. The pairing is described by a BCS approach with a smooth cutoff. For the BCS potential $V(r)$ we have taken a density dependent effective interaction of zero range as introduced by Bertsch and Esbensen[26],

$$V(r) = V_0 [1.0 - \eta (\rho(r)/\rho_0)^\beta], \quad (22)$$

where $\rho_0 = 0.16 \text{ fm}^{-3}$ is the saturation density. Furthermore, the values $V_0 = 481.0$, $\eta = 1.0$, and $\beta = 1.0$ are taken for the parameters V_0 , η , and β . These parameters have been determined by Garrido *et al.*[27].

The anomalous density has the form of

$$\chi(r) = \sum_{nljm} (2j+1) \frac{w_{nljm} u_{nljm} v_{nljm}}{2} \psi_{nljm}(r)^2, \quad (23)$$

where u_{nljm} and v_{nljm} are variational parameters solved from the BCS equations,

$$u_{nljm}^2 = \frac{1}{2} \left[1 + \frac{\epsilon_{nljm} - \epsilon_F}{E_{nljm}} \right]; \quad v_{nljm}^2 = \frac{1}{2} \left[1 - \frac{\epsilon_{nljm} - \epsilon_F}{E_{nljm}} \right] \quad (24)$$

with $E_{nljm} = \sqrt{(\epsilon_{nljm} - \epsilon_F)^2 + \Delta_{nljm}^2}$ the energy of the quasi-particle state and ϵ_{nljm} the single-particle energy determined in the HF equations. Furthermore, $\Psi_{nljm}(\vec{r})$ is the HF orbital wave function of Eq. 15 and w_{nljm} is the smooth cutoff,

$$w_{nljm} = [1.0 + \exp((\epsilon_{nljm} - E_F - E_{cut})/\Delta\epsilon)]^{-1}, \quad (25)$$

with $E_{cut} = 5 \text{ MeV}$ and $\Delta\epsilon = E_{cut}/10$. Therefore, the local gap function can be written as

$$\Delta(r) = V(r)\chi(r), \quad (26)$$

where $V(r)$ is the BCS potential of Eq. 22 and $\chi(r)$ is the anomalous density in Eq. (23).

III. RESULTS AND DISCUSSION

A. Nuclear Matter

All the results which we are going to discuss in this chapter employ a low-momentum interaction V_{lowk} , which is based on the proton-neutron part of CD Bonn potential[8]. Using a cutoff parameter $\Lambda = 2 \text{ fm}^{-1}$, these results do not significantly depend on this choice of the underlying realistic interaction.

First, let us turn to the binding energy of symmetric nuclear matter, which are displayed in Fig. 1. The HF calculation using V_{lowk} does not lead to a minimum in the energy versus density plot (see solid line in Fig. 1), as we have




FIG. 1: (Color online) Binding energy per nucleon of symmetric nuclear matter of a Hartree-Fock calculation using V_{lowk} (solid line) and of a calculation supplemented with a contact interaction (dashed line). Furthermore, comparison is made with a Dirac Brueckner Hartree Fock (DBHF) calculation [32, 33] (dashed-dotted line) and a Brueckner Hartree Fock calculation using the potential Argonne V18 plus a 3-nucleon force (BHF + 3BF) [29] (dotted line).

already mentioned before [13, 16]. The absence of saturation is one of the major problems of calculations for nuclear matter using V_{lowk} . This problem cannot be cured by the inclusion of correlations beyond the HF approximation[15]. Using a conventional model for a realistic interaction, like e.g. the CD Bonn interaction or the Argonne V18, which are not reduced to its low-momentum components, one obtains a saturation point in calculations which account for correlations beyond the mean-field approximation. Therefore one may argue that the V_{lowk} approach cannot reproduce the saturation of nuclear matter as it misses the quenching of short-range correlations in the nuclear medium, which is included in sophisticated calculations using one of the conventional models for a realistic NN interaction.

Note, however, that calculations like Brueckner Hartree Fock (BHF) for these interactions lead to a saturation point, but are not able to reproduce the empirical data. Depending on the amount of tensor and short-range correlations produced by these interactions either the predicted density is too large or the calculated binding energy is too weak[10, 28]. In order to obtain the experimental data for the saturation point one has to include a three-body force[29] or include relativistic effects e.g. within a Dirac Brueckner Hartree Fock (DBHF) approach[30, 31, 32, 33]. The relativistic effects due to the change of the nucleon Dirac spinors in the medium must be considered within the framework of non-relativistic calculations by means of a density-dependent two-nucleon or an effective three-nucleon interaction.

In Fig. 1, our calculation for the binding energy is compared to the ones of a DBHF calculation from Ref. [32, 33] and the non relativistic BHF calculation from Ref. [29]. The non relativistic BHF calculation is based on the Argonne V18 [9] and includes a three-body force deduced from the meson-exchange current approach [29].

Using the V_{lowk} approach, the absence of saturation can be cured by adding three-body forces [13]. Therefore, we also supplemented the V_{lowk} potential with the contact interaction of Eq. (5) to obtain the experimental saturation behavior with a binding energy per nucleon of $E/A = -16.0$ MeV at a saturation density $\rho_0 = 0.16 \text{ fm}^{-3}$. It is worth to notice that only the isoscalar part of the contact interaction can influence the binding energy in symmetric nuclear matter.

In order to fix the isovector part of the contact interaction as well, we also inspect the symmetry energy of infinite matter as a function of density in Fig. 2. The symmetry energy of 21 MeV at saturation density obtained from the V_{lowk} calculation is only about two third of the experimental value. Furthermore, the value of the symmetry energy is significantly below the one predicted by DBHF also at higher densities. This means that the symmetry energy, respectively the isospin dependence of the EoS, of V_{lowk} without contact interaction is in principal too soft.

So we have added a contact interaction term as defined in Eq. (5) choosing a value for $\alpha = 0.5$ and $x_0 = 0.0$. The three remaining parameters have then been fitted to reproduce the empirical saturation point of nuclear matter and the symmetry energy at saturation density. The results for these fitting parameters are listed in Table I and the




FIG. 2: (Color online) The symmetry energy as a function of the density. The symmetry energy of the Hartree-Fock calculations using both V_{lowk} (solid line) and V_{lowk} plus a contact interaction (dashed line) is plotted. For comparison the results of the DBHF approach [33] are plotted as well.

t_0 [MeV fm ³]	x_0	t_3 [MeV fm ^{3+3α}]	x_3
584.1	0.0	8330.7	-0.5

TABLE I: Parameters t_0 , x_0 , t_3 and x_3 defining the contact interaction of Eq. (5)

corresponding results for the binding energy of symmetric nuclear matter and the symmetry energy as a function of density are displayed in Figs. 1 and 2 as well.

The first observation that becomes evident from Fig. 1 is that the V_{lowk} plus contact interaction ($V_{lowk}+CT$) and DBHF calculations yield results which are rather similar, while the result of the BHF approach predicts lower energies at high densities. The compressibility modulus of $V_{lowk}+CT$ at saturation density is $K = 258$ MeV, while those of the other microscopic calculations range from 200 MeV to 265 MeV. This means all the presented equations of state (EoSs) in Fig. 1 can be characterized as rather soft, at least at densities up to about three times saturation density. This prediction of a soft equation of state (EoS) is also supported from observables extracted from heavy ion reactions. For example, heavy ion data for transverse flow [34] or from kaon production [35] support the picture of a soft EoS in symmetric nuclear matter.

In Fig. 2 we can see that the addition of the contact interaction enables us to bring the value of the symmetry energy close to the experimental value. The symmetry energy at saturation density is now 33 MeV. Moreover it is comparable to that of the microscopic DBHF approach up to densities of three times saturation density. Therefore the isospin dependence of the EoS for $V_{lowk}+CT$ calculation is rather stiff.

B. Finite Nuclei

In the first part of this section we are going to consider the bulk properties of nuclei with closed shells for protons and neutrons. The typical example for a heavy nucleus is the example of ^{208}Pb . Results for HF calculations using the V_{lowk} interaction, expanding the single-particle wave functions in a basis of harmonic oscillator (HO) states are presented Fig. 3. The left panel of this figure displays the calculated binding energy as a function of the oscillator length a (see Eq.(21)). The expansion of the single-particle wave functions has been restricted to oscillator states with $N \leq 8$. One finds that the resulting energy is rather sensitive to the choice of the oscillator parameter and leads to very attractive values at small values of a . This gain in binding energy is accompanied by a decrease of the calculated radius of the neutron distribution as can be seen from the right panel in this figure. The nucleus is collapsing to a




FIG. 3: (Color online) Results of Hartree Fock calculations for the energy per nucleon (left panel) and the radius of the neutron distribution (right panel) of ^{208}Pb . The calculations use either the bare V_{lowk} or the V_{lowk} plus contact (CT) interaction. The single particle wave functions are expanded in an oscillator basis. Results are presented as a function of the oscillator length parameter a .




FIG. 4: (Color online) Results of Hartree Fock calculations for the energy per nucleon of ^{16}O . The calculations use either the bare V_{lowk} or the V_{lowk} plus contact (CT) interaction. The single particle wave functions are expanded in an oscillator basis. Results are presented as a function of the oscillator length parameter a .

tightly bound system with a large density. So these results reflect the properties of infinite matter if the bare V_{lowk} interaction is employed.

The situation is rather different if we consider light nuclei. As an example we focus our attention on ^{16}O and present in Fig. 4 the results for the energy per nucleon using various values for the oscillator parameter a . Using V_{lowk} one obtains results which are rather insensitive on the choice of a over a large range of values. Only a basis with a larger than 2 fm seems not to be adequate for such calculation and leads to smaller values for the binding energy per nucleon. Also it is worth noting that the calculated energies around -7.7 MeV per nucleon are rather close to -7.98 MeV observed in experiment. This rather good description of light nuclei using V_{lowk} in an appropriate HO basis has been observed before by Coraggio *et al.*[18, 36].




FIG. 5: (Color online) Comparison of total density distributions calculated for ^{16}O using plane wave basis in a spherical box of radius $R = 15$ fm (PW, solid lines) and an appropriate harmonic oscillator (HO, dashed lines) basis. Results of HF V_{lowk} calculation without contact term (dark lines) and with contact term (light lines) are shown.

How are these results for finite nuclei modified by the contact interaction, which we have added to describe the properties of infinite matter? This is visualized in Figs. 3 and 4 as well. In the case of ^{208}Pb the contact interaction provides a stabilization of the results leading to values for the energy per nucleon and radius of the nucleon distribution, which are in good agreement with the experimental data.

The same is true as well for the case of ^{16}O . The results obtained for the energy per nucleon in HF calculation is very insensitive to the oscillator parameter considered and the calculated energy is rather close to the experimental value.

The calculated energies for ^{16}O with and without the contact interaction are rather close to each other (see Fig. 4), the resulting wave functions, however, are quite different. This can be seen from Fig. 5 showing the calculated nuclear density profiles. The bare V_{lowk} interaction yields densities in the center of the nucleus, which are close to three times the saturation density of nuclear matter. Consequently, the calculated radii are very much below the empirical result. This is in line with the HF calculations of [36]. This also explains why oscillator parameters a larger than 2 fm, are not suitable to describe such a solution. The density profile derived from the $V_{lowk} + \text{CT}$ interaction model yields densities below the saturation density and consequently results for the radius of the charge distribution, which are in good agreement with experiment (see discussion below).

Up to this point we have restricted the discussion to HF calculations performed within an oscillator basis. How do these results compare with the ones derived within the PW basis. Before we proceed to this comparison it should be mentioned that the calculations within the HO basis are much easier to perform than those within the PW basis. Not only that the techniques to calculate the two-body matrix elements (in particular the transformation from relative to laboratory coordinates) is more tediously, also the number of basis states which have to be considered is typically larger in the PW basis as compared to the HO basis. In order to obtain results which are stable from the numerical point of view, we have typically considered a reference box with a radius R which is about 5 times as large as the radius of the nucleus considered. In the case of ^{16}O we have typically considered boxes with radii $R = 15$ fm, while a calculation of ^{208}Pb seems to require a box as large as $R = 30$ fm. This seems to be necessary in order to obtain a set of radial momenta (k_{il} in Eq.(17)), which has a sufficient resolution. For small values of the reference box we also run into the problem that one is considering interactions between neighbored Wigner Seitz cells. For such large

Nucleus	E/A [MeV]			$r_{rms,p}$ [fm]		$r_{rms,n}$ [fm]	
	HO	PW	exp.	HO	PW	HO	PW
^{16}O	-7.77	-7.91	-7.98	2.68	2.68	2.66	2.66
^{40}Ca	-8.33	-8.57	-8.55	3.41	3.41	3.37	3.37
^{48}Ca	-8.13	-8.42	-8.67	3.46	3.45	3.68	3.68
^{60}Ca	-7.42	-7.75	-	3.60	3.59	4.11	4.15
^{208}Pb	-7.25	-7.76	-7.87	5.48	5.45	5.76	5.70

TABLE II: Results for the binding energies per nucleon and rms radii for different nuclei using the V_{lowk} plus contact term interaction model using the harmonic oscillator (HO) or plane wave (PW) basis. The calculated energies have been corrected by subtracting the spurious energy of the center of mass motion. The experimental values are taken from Ref. [37].

^{16}O						
Orbital	Neutron			Proton		
	HO	PW	exp.	HO	PW	exp.
$s_{1/2}$	-37.024	-37.162	-47	-33.478	-33.601	-44 $\pm$ 7
$p_{3/2}$	-19.730	-20.006	-21.839	-16.358	-16.632	-18.451
$p_{1/2}$	-16.217	-16.484	-15.663	-12.888	-13.155	-12.127
$d_{5/2}$	-3.509	-3.739	-4.144	-0.444	-0.690	-0.601
$1s_{1/2}$	-1.288	-1.566	-3.273	1.436	0.839	-0.106
$d_{3/2}$	0.720	0.339	0.941	3.562	1.886	4.399

TABLE III: Single particle energies for the orbital levels of ^{16}O nucleus derived from HF calculations using the V_{lowk} plus contact interaction. Results are displayed using the harmonic oscillator (HO) and plane wave (PW) basis. The experimental values are taken from Ref. [36].

boxes, however, we need a large number of such momenta to cover the relevant range of momenta. This number is typically of the order 15 or larger. The number of different oscillator states, which are required to obtain stable results is significantly smaller.

Comparing the results displayed by dashed and dotted lines in Fig. 5 one find that the results for the density distributions, which are obtained on the PW basis (solid lines) are very close to the ones derived in the HF approximation using the HO basis. This is particularly true for the calculations including the contact interaction term. Note that the HF calculation using the PW basis leads to an energy per nucleon of -7.91 MeV, which is slightly more attractive than the corresponding number of -7.77 MeV obtained in the HO basis. This indicates that the larger number of variational parameters, which are optimized in the PW basis (see discussion above) leads to deeper minimum than the HF variational procedure in the HO basis. Both results are rather close to each other and to the experimental value (see results listed in table II).

This table exhibits very similar findings for the HF calculations of other nuclei with closed shells for protons and neutrons including examples from light (^{16}O) to heavy nuclei (^{208}Pb). It seems that our fitting procedure for the contact term in infinite matter yields very good results for the energies and radii of finite nuclei.

Also the single-particle energies are very similar using the HO or PW basis. This can be seen from table III, which lists the single-particle energies for the example of ^{16}O for the occupied 0s and 0p shells as well as for the 1s0d shell. The single-particle energies evaluated in the PW basis are slightly more attractive than those derived within the HO approximation. The difference is smaller of the tightly bound 0s states than for the more weakly bound states in the 0p and 1s0d shells.

The shapes of corresponding single-particle wave function displayed in Fig. 6 may exhibit the reason for this behavior. The wave functions resulting from the HO and PW cannot be distinguished in this logarithmic plot for radii up to 7 fm. For larger distance from the center, however, the oscillator model yields a faster decrease reflecting the Gaussian asymptotic, which is typical for the HO model.

The differences in the asymptotic behavior of the wave functions are larger, if we consider the shapes of the most weakly bound orbits for protons and neutrons in the case of the weakly bound ^{60}Ca , which are shown in Fig. 7. This is in line with our expectations that the PW basis is superior to the HO basis in particular for the description of weakly bound nucleons.

Finally, we want to study the effect of the basis to be used on the predictions for the occurrence of the neutron drip line. For that purpose we perform HF plus BCS calculations, as described in section II, using the HO as well as the PW basis. As an example we present the results for Fermi energy of neutrons for the Ne isotopes ranging from




FIG. 6: The neutron wave functions for the state $p_{1/2}$ states of ^{16}O calculated in the HF approximation using two different basis systems: the plane wave basis (solid lines) and the harmonic oscillator basis (dashed lines).

$A = 20$ to $A = 32$ in Fig. 8. One can see that the differences in the predictions are rather small for the tightly bound nuclei up to ^{24}Ne and is getting larger with increasing number of neutrons. In this example the oscillator model would predict stable isotopes up to ^{28}Ne , while the calculation using the PW basis predicts stable isotopes up to ^{30}Ne .

IV. SUMMARY AND CONCLUSION

It has been the aim of this study to establish a tool, which allows a microscopic description of nuclei closed to the neutron drip and nuclear structures in the transition from isolated nuclei to homogeneous matter based on a realistic NN interaction. A model space technique, the unitary-model-operator approach (UMOA) [19], has been used to disentangle the low-momentum parts from the high-momentum parts from the CD Bonn interaction[8]. This leads to a universal effective interaction V_{lowk} , which is essentially independent on the underlying realistic interaction V , if the cutoff Λ is chosen around $\Lambda = 2 \text{ fm}^{-1}$.

The Hartree-Fock (HF) calculations using V_{lowk} do not give any saturation in nuclear matter and also HF calculation for finite nuclei lead to nuclear systems of high density and large binding energies. It has been suggested to compensate this feature by adding three-body forces [13]. Instead we supplement the V_{lowk} interaction by a density-dependent contact interaction and adjust the parameters in such a way that HF calculations for infinite matter reproduce the empirical saturation point for symmetric matter and the symmetry energy.

This interaction model is than used to evaluate the properties of finite nuclei. For nuclei with closed shells for protons and neutrons we obtain a very good agreement with the empirical data. Special attention is paid to the technique for solving the HF equations for finite nuclei. The standard technique, expanding the single-particle wave functions in a basis of appropriate harmonic oscillator states is compared to a technique in which these states are expanded in a basis of plane waves. Rather close agreement between these approximation schemes is obtained for nuclei, which are tightly bound. The plane wave basis is favorable, however, for calculating loosely bound systems, since it allows for a better description of the asymptotic part of the nucleon wave functions.

These tools, the interaction model and the technique for solving the HF equations, can now be used to describe the structure of loosely bound nuclei, like halo nuclei or nuclei at the edges of the valley of stability, or the nuclear structures in the so-called pasta phase of neutron stars. Another perspective is to consider a many-body approach




FIG. 7: (Color online) The neutron wave function of $1p_{1/2}$ (dark lines) and the proton wave function of $d_{3/2}$ of ^{60}Ca (light lines) using two different basis systems, the plane wave basis (solid lines) and the harmonic oscillator basis (dashed lines).




FIG. 8: (Color online) The Fermi energies for the neutrons in the isotopes of Ne . Results of HF plus BCS calculations using the plane wave basis (PW) and the harmonic oscillator basis (HO) are shown for various even numbers of neutrons.

for describing the nuclear systems, which goes beyond the HF approximation.

V. ACKNOWLEDGEMENTS

This work has been supported by a grant (Mu 705/5-1) of the Deutsche Forschungsgemeinschaft DFG

-
- [1] T.H.R. Skyrme, Nucl. Phys. **9**, 615 (1959).
 - [2] D. Vautherin and D.M. Brink, Phys. Rev. C **5**, 626, (1972).
 - [3] P. Bonche and D. Vautherin, Nucl. Phys. A **372**, 496 (1981).
 - [4] K. Oyamatsu and K. Iida, Phys. Rev. C **75**, 015801 (2007).
 - [5] F. Montani, C. May, and H. M  ther, Phys. Rev. C **69**, 065801 (2004).
 - [6] P. G  gelein and H. M  ther, Phys. Rev. C **77**, 024312 (2008).
 - [7] P. G  gelein, E. N. E. van Dalen, C. Fuchs, and H. M  ther, Phys. Rev. C **77**, 025802 (2008).
 - [8] R. Machleidt, F. Sammarruca, and Y. Song, Phys. Rev. C **53**, R1483 (1996).
 - [9] R.B. Wiringa, V.G.J. Stoks, and R. Schiavilla, Phys. Rev. C **51**, 38 (1995).
 - [10] H. M  ther and A. Polls, Prog. Part. Nucl. Phys. **45**, 243 (2000).
 - [11] S.C. Pieper, R.B. Wiringa, and J. Carlson, Phys. Rev. C **70**, 054325 (2006).
 - [12] S. K. Bogner, T.T.S. Kuo, and L. Coraggio, Nucl. Phys. A **684**, 432c (2001).
 - [13] S.K. Bogner, A. Schwenk, R.J. Furnstahl, and A. Nogga, Nucl. Phys. A **763**, 59 (2005).
 - [14] S. K. Bogner, R. J. Furnstahl, S. Ramanan, and A. Schwenk, Nucl. Phys. A **784**, 79 (2007).
 - [15] P. Bo  zek, D.J. Dean, and H. M  ther, Phys. Rev. C **74**, 014303 (2006).
 - [16] J. Kuckei, F. Montani, H. M  ther, and A. Sedrakian, Nucl. Phys. A **723**, 32 (2003).
 - [17] L. Coraggio, A. Covello, A. Gargano, N. Itaco, T. T. S. Kuo, and R. Machleidt, Phys. Rev. C **71**, 014307 (2005).
 - [18] L. Coraggio, A. Covello, A. Gargano, N. Itaco, and T. T. S. Kuo, Phys. Rev. C **75**, 057303 (2007); Phys. Rev. C **73**, 014304 (2006).
 - [19] K. Suzuki, Prog. Theoret. Phys. **68**, 246 (1986).
 - [20] S. Fujii, R. Okamoto, and K. Suzuki, Phys. Rev. C **69**, 034328 (2004).
 - [21] M. Moshinsky, Nucl. Phys. **13**, 104 (1959).
 - [22] I. Talmi, Helv. Phys. Acta **25**, 185 (1952).
 - [23] I. Talmi, “Simple Models of Complex Nuclei”, Harwood Academic Publishers (Chur, Switzerland, 1993).
 - [24] C.L. Kung, T.T.S. Kuo, and K.F. Ratcliff, Phys. Rev. C **19**, 1063 (1979).
 - [25] D. Bonatsos and H. M  ther, Nucl. Phys. A **496**, 23 (1989).
 - [26] G.F. Bertsch and H. Esbensen, Ann. Phys. **209**, 327 (1991).
 - [27] E. Garrido, P. Sarriguren, E. Moya de Guerra, and P. Schuck, Phys. Rev. C **60**, 064312 (1999).
 - [28] F. Coester, S. Cohen, B.D. Day, and C.M. Vincent, Phys. Rev. C **1**, 769 (1970).
 - [29] A. Lejeune, U. Lombardo, and W. Zuo, Phys.Lett. B **477**, 45 (2000).
 - [30] R. Brockmann and R. Machleidt, Phys. Lett. B **149**, 283 (1984).
 - [31] H. M  ther, R. Machleidt, and R. Brockmann, Phys. Rev. C **42**, 1981 (1990).
 - [32] E. N. E. van Dalen, C. Fuchs, and Amand Faessler, Nucl. Phys. A **744**, 227 (2004).
 - [33] E.N.E. van Dalen, C. Fuchs, and A. Faessler, Eur. Phys. J. A **31**, 29 (2007).
 - [34] G. Stoicea *et al.* [FOPI Coll.], Phys. Rev. Lett. **92**, 072303 (2004).
 - [35] C. Sturm *et al.* [KaoS Coll.], Phys. Rev. Lett. **86**, 39 (2001); C. Fuchs, A. Faessler, E. Zabrodin, Y.M. Zheng, Phys. Rev. Lett. **86**, 1974 (2001); C. Fuchs, Prog. Part. Nucl. Phys. **56**, 1 (2006).
 - [36] L. Coraggio, N. Itaco, A. Covello, A. Gargano, and T. T. S. Kuo, Phys. Rev. C **68**, 034320 (2003).
 - [37] F. Hofmann, C.M. Keil, and H. Lenske, Phys. Rev. C **64**, 034314 (2001).